

\_\_\_\_\_

ProM1 <sup>™</sup> MHC Class I Monomer (Biotin-labeled):	MHC Class I-peptide complexes bind to CD8+ T cell receptors of a particular specificity, as determined by the MHC allele and peptide combination. In addition to their use (as streptavidin-conjugated MHC tetramers) in the identification of specific CD8+ T cells by flow cytometry, biotin-labeled MHC Class I monomers can be used to isolate (or deplete) antigen specific CD8+ T-cells through the use of streptavidin-coated magnetic microbeads. Isolation of antigen-specific T cells in this manner is useful if viable cells are needed for downstream applications. Biotin-labeled MHC Class I monomers can also be immobilised on streptavidin-coated surfaces for use in plate based assays such as ELISA. <b>For Research Use Only. Not for use in therapeutic or diagnostic procedures</b> .
Test size:	ProM1 <sup>™</sup> Monomers are provided in 35 µg and 100 µg sizes
Concentration/ Formulation:	ProM1 <sup>™</sup> Monomers are supplied at a concentration of approximately 0.4 mg/mL in PBS, stabilized with 1% BSA and 0.025% sodium azide.
Storage Condition:	Use liquid nitrogen to flash-freeze upon receipt and store at -80 °C. Avoid freeze-thaw cycles.
Shelf Life:	The monomer is stable for 12 months if stored as instructed above.
Hazards:	This reagent is formulated in 0.025% sodium azide. Under acidic conditions the toxic compound hydrazoic acid may be released. Compounds containing sodium azide should be flushed with running water while being discarded.

## Quality Control Assay Results

Appearance: Colorless solution

Protein Characterization: Passed

**Released by:** (Date as per product label above)

© Copyright ProImmune Limited 2003-2021. All Rights Reserved. PS\_M (ProM1 MHC Class I Monomer) Version 1.1

Page 1 of 1